






Mulai vang baru














Scientific Foundation SPIROSKI, Skopje, Republic of Macedonia Open Access Macedonian Journal of Medical Sciences. 2021 Feb 05; 9(B):1-3. https://doi.org/10.3889/oamjms.2021.7495 eISSN: 1857-9655

Category: B - Clinical Sciences Section: Infective Diseases



AQ3

AQ2

AQ4

AQ5

AQ2



# Synthetic Food Coloring as an Alternative to Worm Egg Staining Ascaris lumbricoides

AQ2,7 Steven Jacub Soenjono<sup>1</sup>, Budi Setiawan<sup>2,3</sup>, Rahmah Siti Wulandari<sup>2</sup>, Suyana<sup>2</sup>, Marta Atik Martiningsih<sup>2</sup>

<sup>1</sup>Department of Environmental Health, Politeknik Kesehatan Kementerian Kesehatan Manado, Indonesia; <sup>2</sup>Department of Medical Laboratory Technology, Politeknik Kesehatan Kementerian Kesehatan Yogyakarta, Indonesia; <sup>3</sup>Department of ???, Pusat Unggulan IPTEKS Inovasi Teknologi Kesehatan Masyarakat (PUINOVAKESMAS), Politekkes Kemenkes Yogyakarta, Indonesia

### Abstract

Edited by: https://publons.com/researcher/391987/
mirko-spiroski
Citation: Soenjono SJ, Setiawan B, Wulandan RS,
Suyana, Martiningsih MA. Synthetic Food Coloring as an
Alternative to Worm Egg Staining Ascaris lumbricoides.
Open Access Maced J Med Sci. 2021 Feb. 05; 9(B):1-3.
https://doi.org/10.3889/oamjms.2021.7495
Keywords: Ascaris lumbricoides; Carmoisin Cl 14720.
Eosin 2%

Eosin 2%
\*Correspondence: ???
Received: ???
Revised: ???

Copyright: © 2021 Steven Jacub Soenjono,
Budi Setiawan, Rahmah Siti Wulandari, Suyana,
Marta Alti Martiningsih
Funding: This research did not receive any financial
support
Competing Interest: The authors have declared that no
competing interest exists
Open Access: This is an open-access article distributed
under the terms of the Creative Commons AttributionNonCommercial 4.0 International License (CC BY-NC 4.0)

Examination of worm eggs is a way of diagnosing helminthiasis by giving 2% eosin as a dye, but this reagent is carcinogenic so an alternative is needed to replace it. Synthetic food coloring (carmoisin Cl 14720) can staining worm eggs against a more contrasting background than eosin. The purpose of this study was to determine the synthetic food coloring with a concentration of 1%, 2%, 3%, 4%, and 5% which can be used as an alternative to 2% eosin in the examination of *Ascaris lumbricoides* eggs. This type of research is observational with a cross-sectional research design. Positive stool samples for *A lumbricoides* eggs were obtained from the Parasitology Laboratory of the University of Indonesia. Data were analyzed by ANOVA and interclass correlation coefficient (ICC) tests. The results of the ANOVA test showed that there was no difference between 2% eosin and food coloring. The results of the ICC test obtained a value of 0.845, so the dye can be used as an alternative to 2% eosin in the examination of *A. lumbricoides* eggs.

### Introduction

One-third of the world's population is infected with at least one species of soil-transmitted helminths (STHs) with Ascaris lumbricoides infecting 800 million people, Trichuris trichiura 600 million, and hookworm 600 million and causing up to 135,000 deaths annually [1]. STHs have also been classified among the most common neglected tropical diseases because they persist exclusively in the poorest populations (i.e., US\$1,25/day) and are thus often overlooked [2]. Individuals infected with STH may suffer from anemia, stunted growth, reduced physical fitness, and impaired cognitive development [3] which show a continued decline in socioeconomic development in low-income countries [4], [5].

At present, the identification of parasitic worms is more focused on examining eggs as the basis for determining the genus of worms, although it is generally known that many parasitic eggs are relatively the same [6]. Identification can be done in several ways, one of which is by direct (native) examination of the preparation. The dye reagent used in this examination is 2% eosin which is acidic, which can provide a red background and produce a red morula color and dark

red wall color in worm eggs [7]. However, it is suspected that eosin can be a carcinogen in living things [8], therefore, alternative materials are needed to replace them.

One of the dyes that are easily available is synthetic dyes for food. The dye contains carmoisin dye with CI code 144,720 which produces a dark red color. Carmoisin is a synthetic dye (water soluble) containing an azo structure and an aromatic ring [9] and not carcinogenic [10]. Many researches related to alternative materials to replace eosin have been carried out, but there has been no research using carmoisin to color *A. lumbricoides* eggs. In this study, we evaluated the quality of carmoisin as a potential alternative for staining *A. lumbricoides* compared to 2% eosin.

# **Methods**

This research is observational analytic with cross-sectional study, feces examined with direct method. Samples of *A. lumbricoides* eggs were obtained from the Parasitology Laboratory of

1

B - Clinical Sciences Infective Diseases

the University of Indonesia. The fecal sample was examined by 10 examiner to give a score with the criteria of a score of 1, the preparation had a noncontrast field of view, the worm eggs did not absorb color and the egg part was not clear, the score was 2, if the preparation had a less contrasting field of view, the worm eggs did not absorb color, and the eggs are not clearly visible, the score is 3 when the visual field is contrasting, the worm eggs absorb the color and the egg parts are clearly visible.

The data are presented in a table of each concentration then ANOVA analysis is carried out then followed by interclass correlation coefficient (ICC) analysis to determine the ICC value at each concentration to determine the reliability of the examination results.

### **Results and Discussion**

Based on the research results shown in Figure 1, the food coloring used can color all parts of the egg, but the best concentration is at 2%. Research conducted by Nurmufidah (2020), in the description of the variation in the concentration of Giemsa solution on the quality of STH worm egg preparations, it was found that the quality of preparations with the best quality at a concentration of 3% with a constant white background. easier, and more comfortable to distinguish with worm eggs, the embryo is purplish with a purplish-blue color on the egg wall of the Necator americanus/Ancylostoma duodenale with 150% effectiveness so that it can be used as a substitute for 2% eosin in the manufacture of worm egg preparations by the direct method. While this study used a synthetic food coloring carmoisin CI 14720 with the results of the preparation that is a white background with pink eggs, morula can be seen clearly.




Figure 1: Microscopic view of staining worm eggs

The principle of staining on worm eggs is the affinity (formation of chemical bonds) between the tissue and the dye, either directly or indirectly [11]. Carmoisin CI 14720 dye tends to be slightly alkaline [12]. This is evidenced by the examination of the pH of carmoisin CI 44720 at each concentration having the same pH of 7.5. Carmoisin CI 14720 contains an azo structure and an aromatic ring [13] Carmoisin uses propylene glycol as a solvent. Propylene glycol functions as a humectant that will maintain the stability of the preparation by

Table 1: Carmoisin staining analysis

Examiner	Carmoisin dye concentration					Eosin	ICC
	1%	2%	3%	4%	5%	2%	-
1	1	3	3	2	3	2	0.845
2	2	3	3	1	2	3	
3	3	3	3	2	2	3	
4	3	2	3	2	2	3	
5	3	3	3	2	2	3	
6	3	3	2	2	2	2	
7	3	3	3	3	3	3	
8	3	3	3	3	3	3	
9	3	3	2	1	2	3	
10	3	3	3	2	2	3	
ANOVA test	0.102	0.645	0.286	0.242	0.486		

reducing evaporation of water from the preparation and absorbing moisture from the environment [14].

Worm eggs contain albuminoid which is a type of protein [15]. Protein molecules have an amino group (-NH2) and a carboxylic group (-COOH) at the ends of the chain [16]. This causes the protein to have a lot of charge (polyelectrolyte) and is amphoteric, that is, it can react with acids and bases [17] then the chemical bond formed between carmoisin and protein is a carboxylic group reacting with OH- ions, the presence of a charge on the protein molecule causes the protein to move under the influence of an electric field [18] so that the negatively charged protein that causes *A. lumbricoides* worm eggs can be stained by positively charged carmoisin CI 14720.

### Conclusion

Carmoisin CI 14720 synthetic food coloring can be used as an alternative dye for *A. lumbricoides* worm eggs with an ICC value of 0.845. Further research can examine the effect of the synthetic food coloring carmoisin CI 14720 on the examination of worm eggs with different worm eggs.

### References

- Hotez PJ. One world health: Neglected tropical diseases in a flat world. PLoS Negl Trop Dis. 2009;3(4):e405. http://doi. org/10.1371/journal.pntd.0000405
  - PMid:19399165
- Liese B, Rosenberg M, Schratz A. Programmes, partnerships, and governance for elimination and control of neglected tropical diseases. Lancet. 2010;375(9708):67-76. http://doi.org/10.1016/ S0140-6736(09)61749-9
  - PMid:20109865
- Bethony J, Brooker S, Albonico M, Geiger SM, Loukas A, Diemert D, et al. Soil-transmitted helminth infections: Ascariasis, trichuriasis, and hookworm. Lancet. 2006;367(9521):1521-32. http://doi.org/10.1016/S0140-6736(06)68653-4
  - PMid:16679166
- 4. King CH. Parasites and poverty: The case of schistosomiasis.

Acta Trop. 2010;113(2):95-104. http://doi.org/10.1016/j.actatropica.2009.11.012

PMid:19962954

- World Health Organization. Deworming for Health and Development: Report of the 3<sup>rd</sup> Global Meeting of the Partners for Parasite Control. Geneva: World Health Organization; 2005.
- Hanafiah M, Dwinna A, Mahdi A, Karmil F, Rachmady D. Detection of parasitic helminths in cattle from Banda Aceh, Indonesia. Vet World. 2019;12(8):1175. http://doi.org/10.14202/ vetworld.2019.1175-1179

PMid:31641294

- Oktari A, Mu'tamir A. Optimasi air perasan buah merah (*Pandanus* sp.) pada pemeriksaan telur cacing. J Teknol Lab. 2017;6(1):8-17.
- Adekunle AA. Studies of age-related changes in the metabolism of eosin in the rat. Mech Ageing Dev. 1978;8(3):219-25. http:// doi.org/10.1016/0047-6374(78)90020-9
   PMid:211356
- Oveisi M, Sadeghi N, Jannat B, Nilfroush E. Simultaneous determination of carmoisine and ponceau 4R. Food Anal Methods. 2008;1(3):214-9. http://doi.org/10.1007/

s12161-008-9022-7

 Mason PL, Gaunt IF, Butterworth KR, Hardy J, Kiss IS, Grasso P. Long-term toxicity studies of carmoisine in mice. Food Cosmet Toxicol. 1974;12(5-6):601-7. http://doi. org/10.1016/0015-6264(74)90232-6
 PMid:4452535

- 11. Jusuf AA. Histoteknik Dasar. Indonesia: Universitas Indonesia; 2009.
- WC Analisis dan Aspek Kesehatan Bahan Tambahan Pangan. Bumi Aksara: 2006.
- 13. Sumarlin LO. Identifikasi Pewarna Sintetik Pada Produk Pangan Yang Beredar di Jakarta dan Ciputat; 2010.
- Martin A, Swarbick J, Cammarata A. Farmasi Fisik 2. UI Press; 1993.
- 15. Poedjiadi A. Dasar-Dasar Biokimia. Ul Press; 1994.
- 16. Winarno. Kimia Pangan dan Gizi. Gramedia; 2008.
- 17. Yusuf Y. Modul Sederhana dan Ilmiah Untuk BELAJAR KIMIA Pangan dan Gizi. Indonesia: EduCenter Indonesia; 2018.
- Yazid EN, Lisda. Penuntun Praktikum Biokimia Untuk Mahasiswa Analis. Andi Offset; 2006.

# Author Queries???

AQ1: Kindly provide running title.
AQ2: Kindly provide author full name.
AQ3: Kindly provide department.

AQ4: Kindly provide corresponding author details.

AQ5: Kindly provide history details AQ6: Kindly cite table 1 in text part.

AQ7: Kindly provide ORCID Id for all authors



Scientific Foundation SPIROSKI, Skopje, Republic of Macedonia Open Access Macedonian Journal of Medical Sciences. 2021 Feb 05; 9(B):1-3. https://doi.org/10.3889/oamjms.2021.7495 elSSN: 1857-9655

Category: B - Clinical Sciences Section: Infective Diseases





# Synthetic Food Coloring as an Alternative to Worm Egg Staining Ascaris lumbricoides

Steven Jacob Soenjono<sup>1</sup>, Budi wan<sup>2,3</sup>, Rahmah S ulandari<sup>2</sup>, S a<sup>2</sup>, Marta Atik Martiningsih<sup>2</sup> AQ2,7

<sup>1</sup>Department of Environmental Health, Politeknik Kesehatan Kementerian Kesehatan Mana<mark>do</mark>, Indones<mark>ia,</mark> <sup>2</sup>Department of Medical Laboratory Technology, Politeknik Kesehatan Kementerian Kesehatan Yogyakarta, Indonesia; <sup>3</sup>Department of Unggulan IPTEKS Inovasi Teknologi Kesehatan Masyarakat (PUINOVAKESMAS), Poltekkes Kemenkes Yogyakarta, Indonesia

### Abstract

Examination of worm eggs is a way of diagnosing helminthiasis by giving 2% eosin as a dye, but this reagent is carcinogenic so an alternative is needed to replace it. Synthetic food coloring (carmoisin CI 14720) can staining worm eggs against a more contrasting background than eosin. The purpose of this study was to determine the synthetic food coloring with a concentration of 1%, 2%, 3%, 4%, and 5% which can be used as an alternative to 2% eosin in the examination of Ascaris lumbricoldes eggs. This type of research is observational with a cross-sectional research design. Positive stool samples for A lumbricoides eggs were obtained from the Parasitology Laboratory of the University of Indonesia. Data were analyzed by ANOVA and interclass correlation coefficient (ICC) tests. The results of the ANOVA test showed that there was no difference between 2% eosin and food coloring. The results of the ICC test obtained a value of 0.845, so the dye can be used as an alternative to 2% eosin in the examination of A. lumbricoides eaas.

AQ2

AQ4

AQ5

AQ2

ation: Soeniono S.J. Setiawan B. Wulandari RS Mation: Soenjono SJ, Setiawan B, Wulandan RS, urap Matriningsih MA. Synthetic Food Coloring as an ative to Worm Egg Staining Ascaris lumbricoides. Open Access Maced J Med Sci. 2021 Feb 05; 9(B):1-3. https://doi.org/10.3889/oamjins.2021.7495 Keywords: Ascaris lumbricoides; Carmoisin Cl 147

Edited by: https://publons.com/researcher/391987/



Competing Interest: The authors have declared that no

Open Access: This is an open-access article distributed under the terms of the Creative Commons Attribution-NonCommercial 4.0 International License (CC BY-NC 4.0)

### Introduction

One-third of the world's population is infected with at least one species of soil-transmitted helminths (STHs) with Ascaris lumbricoides infecting 800 million people, Trichuris trichiura 600 million, and hookworm 600 million and causing up to 135,000 deaths annually [1]. STHs have also been classified among the most common neglected tropical diseases because they persist exclusively in the poorest populations (i.e., US\$1.25/day) and are thus often overlooked [2]. Individuals infected with STH may suffer from anemia, stunted growth, reduced physical fitness, and impaired cognitive development [3] which show a continued decline in socioeconomic development in low-income countries [4], [5].

At present, the identification of parasitic worms is more focused on examining eggs as the basis for determining the genus of worms, although it is generally known that many parasitic eggs are relatively the same [6]. Identification can be done in several ways, one of which is by direct (native) examination of the preparation. The dye reagent used in this examination is 2% eosin which is acidic, which can provide a red background and produce a red morula color and dark red wall color in worm eggs [7]. However, it is suspected that eosin can be a carcinogen in living things [8], therefore, alternative materials are needed to replace them.

One of the dyes that are easily available is synthetic dyes for food. The dye contains carmoisin dye with CI code 144,720 which produces a dark red color. Carmoisin is a synthetic dye (water soluble) containing an azo structure and an aromatic ring [9] and not carcinogenic [10]. Many researches related to alternative materials to replace eosin have been carried out, but there has been no research using carmoisin to color A. lumbricoides eggs. In this study, we evaluated the quality of carmoisin as a potential alternative for staining A. lumbricoides compared to 2% eosin.

# **Methods**

This research is observational analytic with cross-sectional study, feces examined with direct method. Samples of A. lumbricoides eggs were obtained from the Parasitology Laboratory of

1

the University of Indonesia. The fecal sample was examined by 10 examiner to give a score with the criteria of a score of 1, the preparation had a noncontrast field of view, the worm eggs did not absorb color and the egg part was not clear, the score was 2, if the preparation had a less contrasting field of view, the worm eggs did not absorb color, and the eggs are not clearly visible, the score is 3 when the visual field is contrasting, the worm eggs absorb the color and the egg parts are clearly visible.

The data are presented in a table of each concentration then ANOVA analysis is carried out then followed by interclass correlation coefficient (ICC) analysis to determine the ICC value at each concentration to determine the reliability of the examination results.

### **Results and Discussion**

Based on the research results shown in Figure 1, the food coloring used can color all parts of the egg, but the best concentration is at 2%. Research conducted by Nurmufidah (2020), in the description of the variation in the concentration of Giemsa solution on the quality of STH worm egg preparations, it was found that the quality of preparations with the best quality at a concentration of 3% with a constant white background. easier, and more comfortable to distinguish with worm eggs, the embryo is purplish with a purplish-blue color on the egg wall of the Necator americanus/Ancylostoma duodenale with 150% effectiveness so that it can be used as a substitute for 2% eosin in the manufacture of worm egg preparations by the direct method. While this study used a synthetic food coloring carmoisin CI 14720 with the results of the preparation that is a white background with pink eggs, morula can be seen clearly.




Figure 1: Microscopic view of staining worm eggs

The principle of staining on worm eggs is the affinity (formation of chemical bonds) between the tissue and the dye, either directly or indirectly [11]. Carmoisin CI 14720 dye tends to be slightly alkaline [12]. This is evidenced by the examination of the pH of carmoisin CI 44720 at each concentration having the same pH of 7.5. Carmoisin CI 14720 contains an azo structure and an aromatic ring [13] Carmoisin uses propylene glycol as a solvent. Propylene glycol functions as a humectant that will maintain the stability of the preparation by



### e 1: Carmoisin staining analysis

		A
sin	ICC	
)		
	0.845	

Examiner	Carmoisin dye concentration					Eosin	ICC
	1%	2%	3%	4%	5%	2%	-
1	1	3	3	2	3	2	0.845
2	2	3	3	1	2	3	
3	3	3	3	2	2	3	
4	3	2	3	2	2	3	
5	3	3	3	2	2	3	
6	3	3	2	2	2	2	
7	3	3	3	3	3	3	
8	3	3	3	3	3	3	
9	3	3	2	1	2	3	
10	3	3	3	2	2	3	
ANOVA test	0.102	0.645	0.286	0.242	0.486		

reducing evaporation of water from the preparation and absorbing moisture from the environment [14].

Worm eggs contain albuminoid which is a type of protein [15]. Protein molecules have an amino group (-NH2) and a carboxylic group (-COOH) at the ends of the chain [16]. This causes the protein to have a lot of charge (polyelectrolyte) and is amphoteric, that is, it can react with acids and bases [17] then the chemical bond formed between carmoisin and protein is a carboxylic group reacting with OH- ions, the presence of a charge on the protein molecule causes the protein to move under the influence of an electric field [18] so that the negatively charged protein that causes A. *lumbricoides* worm eggs can be stained by positively charged carmoisin CI 14720.

### Conclusion

Carmoisin CI 14720 synthetic food coloring can be used as an alternative dye for A. lumbricoides worm eggs with an ICC value of 0.845. Further research can examine the effect of the synthetic food coloring carmoisin CI 14720 on the examination of worm eggs with different worm eggs.

### References

- Hotez PJ. One world health: Neglected tropical diseases in a flat world. PLoS Negl Trop Dis. 2009;3(4):e405. http://doi. org/10.1371/journal.pntd.0000405
  - PMid:19399165
- Liese B, Rosenberg M, Schratz A. Programmes, partnerships, and governance for elimination and control of neglected tropical diseases. Lancet. 2010;375(9708):67-76. http://doi.org/10.1016/ S0140-6736(09)61749-9
  - PMid:20109865
- Bethony J, Brooker S, Albonico M, Geiger SM, Loukas A, Diemert D. et al. Soil-transmitted helminth infections: Ascariasis. trichuriasis, and hookworm. Lancet. 2006;367(9521):1521-32. http://doi.org/10.1016/S0140-6736(06)68653-4
  - PMid:16679166
- King CH. Parasites and poverty: The case of schistosomiasis.



Acta Trop. 2010;113(2):95-104. http://doi.org/10.1016/j.actatropica.2009.11.012

PMid:19962954

- World Health Organization. Deworming for Health and Development: Report of the 3<sup>rd</sup> Global Meeting of the Partners for Parasite Control. Geneva: World Health Organization; 2005.
- Hanafiah M, Dwinna A, Mahdi A, Karmil F, Rachmady D. Detection of parasitic helminths in cattle from Banda Aceh, Indonesia. Vet World. 2019;12(8):1175. http://doi.org/10.14202/ vetworld.2019.1175-1179

PMid:31641294

- Oktari A, Mu'tamir A. Optimasi air perasan buah merah (*Pandanus* sp.) pada pemeriksaan telur cacing. J Teknol Lab. 2017;6(1):8-17.
- Adekunle AA. Studies of age-related changes in the metabolism of eosin in the rat. Mech Ageing Dev. 1978;8(3):219-25. http:// doi.org/10.1016/0047-6374(78)90020-9
   PMid:211356
- Oveisi M, Sadeghi N, Jannat B, Nilfroush E. Simultaneous determination of carmoisine and ponceau 4R. Food Anal Methods. 2008;1(3):214-9. http://doi.org/10.1007/

s12161-008-9022-7

- Mason PL, Gaunt IF, Butterworth KR, Hardy J, Kiss IS, Grasso P. Long-term toxicity studies of carmoisine in mice. Food Cosmet Toxicol. 1974;12(5-6):601-7. http://doi. org/10.1016/0015-6264(74)90232-6
  - PMid:4452535
- 11. Jusuf AA. Histoteknik Dasar. Indonesia: Universitas Indonesia; 2009.
- WC Analisis dan Aspek Kesehatan Bahan Tambahan Pangan. Bumi Aksara: 2006.
- 13. Sumarlin LO. Identifikasi Pewarna Sintetik Pada Produk Pangan Yang Beredar di Jakarta dan Ciputat; 2010.
- Martin A, Swarbick J, Cammarata A. Farmasi Fisik 2. UI Press; 1993.
- 15. Poedjiadi A. Dasar-Dasar Biokimia. Ul Press; 1994.
- 16. Winarno. Kimia Pangan dan Gizi. Gramedia; 2008.
- 17. Yusuf Y. Modul Sederhana dan Ilmiah Untuk BELAJAR KIMIA Pangan dan Gizi. Indonesia: EduCenter Indonesia; 2018.
- Yazid EN, Lisda. Penuntun Praktikum Biokimia Untuk Mahasiswa Analis. Andi Offset; 2006.

# Author Queries???

AQ1: Kindly provide running title.
AQ2: Kindly provide author full name.
AQ3: Kindly provide department.

AQ4: Kindly provide corresponding author details.

AQ5: Kindly provide history details AQ6: Kindly cite table 1 in text part.

AQ7: Kindly provide ORCID Id for all authors





### **Copyright Form**

Manuscript Number: \_\_7495\_\_

This Agreement is made with *Open Access Macedonian Journal of Medical Sciences (OAMJMS)*. The Agreement is for the Article submitted by you ("Author") for publication *Open Access Macedonian Journal of Medical Sciences (OAMJMS)*. To enable publishing the Article in *Open Access Macedonian Journal of Medical Sciences (OAMJMS)* ("the Journal"), the ownership of copyright must be established. Please read and complete the form below (Section A plus Section B) and return one copy to the address given below. There are explanatory notes to this Agreement, which also form part of the Agreement.

The Article cannot be published until this signed Agreement is received by OAMJMS.

### SECTION A: Author's Warranty (Please print your details)

Name: Steven Jacub Soenjono

Address: Jl. R. W. Mongisidi Malalayang II Kota Manado, Indonesia

Article title: Synthetic Food Coloring as an Alternative to Worm Egg Staining Ascaris lumbricoides

Names of all authors in the order in which they appear in the Article:

- 1. Steven Jacub Soenjono
- 2. Budi Setiawan
- 3. Rahma Siti Wulandari
- 4. Suyana Suyana
- 5. Marta Atik Martiningsih

In consideration of the publication of the Article in the above Journal, I hereby warrant and undertake:

- a. that this Article is an original work, has not been published before and is not being considered for publication elsewhere in its final form either in printed or electronic form.
- b. that I have obtained permission from the copyright holder to reproduce in the Article (in all media including print and electronic form) material not owned by me, and that I have acknowledged the source;
- c. that this Article contains no violation of any existing copyright or other third party right or any material of an obscene, indecent or otherwise unlawful nature and that to the best of my knowledge this Article does not infringe the rights of others;
- d. that I will indemnify and keep indemnified the Editors, Open Access Macedonian Journal of Medical Sciences against all claims and expenses (including legal costs and expenses) arising from any breach of this warranty and the other warranties on my behalf in this Agreement;
- e. that in the case of a multi-authored Article I have obtained copyright assignment from all co-authors, in writing, and authorization to enter into this Agreement on their behalf and that all co-authors have read and agreed the above warranties;

Author: Signed: Steven Jacub Soenjono ...

.. Date: November, 23, 20021

**SECTION B: Copyright Assignment** 

This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited. Notwithstanding the above, I retain all proprietary rights other than copyright, such as patent and trade mark rights and rights to any process or procedure described in the Article.

Author: Signed: Steven Jacub Soenjono .....

. Date: November, 23, 20021

In signing this form, the signee asserts that any authors not signing have authorised the signee to do so on their behalf, and that the manuscript submitted has been approved by these authors in the form in which it has been submitted, and that the warranties given above have been read and agreed by all authors.

new