Halaman Sampul Jurnal



A - Basic Sciences Vol. 9 No. A (2021)

Daftar Isi Artikel





The Effect of Coffee Arabica Gayo Leaf Extract (Coffea arabica L.) in Increasing Phosphoinositide 3-kinase and Glucose Transporter-4 Expression in the Skeletal Muscle

Sake Juli Martina, Aznan Lelo, Dharma Lindarto, Ratna Akbari Ganie, Muhammad Ichwan, Hanifah Yusuf, Syafruddin Ilyas, Iqbal Pahlevi Nasution (Author)

906-910

PDF

The The Prevalence of Potentially Inappropriate Prescribing in the Elderly at the Primary Care Level in Kazakhstan

Ainash Ibysheva, Gulmira Muldaeva, Leila I. Arystan, Almagul B. Kuzgibekova, Bibigul A. Abeuova, Leila S. Haydargalieva 572-578 (Author)

PDF

Alkaloids of Peganum harmala L. and their Pharmacological Activity

Aidos Doskaliyev, Roza Seidakhmetova, D. S. Tutai, Kristina Goldaeva, V.K. Surov, S.M. Adekenov (Author)

766-775

PDF

The Effect of Levamlodipine in Glucose-Induced Acute Model of Glaucoma in Rabbits

Waleed K. Abdulsahib (Author)

505-509

PDF

Evaluation of Antioxidant Activity of Botto-Botto Leaf Fraction (Chromolaena Odorata L.) Using DPPH and ABTS Methods

Karlina Amir Tahir, Upik A. Miskad, Khairuddin Djawad, Sartini Djide, Khaerani Khaerani, Maulita Indrisari (Author)

183-188

PDF

Antibacterial Screening of Endophytic Fungus Xylaria sp. derived from Andrographis paniculata (Sambiloto)

Suryelita Suryelita, Riga Riga, Sri Benti Etika (Author); Mariam Ulfah; Muh Ade Artasasta (Author)

971-975

PDF

The Possible Antidiabetic Effect of Ficus carica L. Tablet on lloxan-Induced Diabetes Model in Rats

Muhammad Fariez Kurniawan, Fadhilah Alvari Yusuf (Author)

727-734

PDF

Effect of Enhydra fluctuans on Kidney Function in Alloxan-induced Diabetic Rats

Rina Delfita, Dahelmi Dahelmi, Djong Tjong, Suhatri Suhatri (Author)

1187-1194

PDF

Formulation and Pharmacological Studies of Leaves of Moringa (Moringa oleifera), a Novel Hepatoprotection in Oral Drug Formulations

Aristianti Aristianti, Nurkhaeri Nurkhaeri, Vanny Y. Tandiarrang, Akbar Awaluddin, Lukman Muslimin (Author)

151-156

PDF

The Effect of Tiwari Onion (Eleutherine americana Merr) Tablet on Blood Pressure Stability in Diagnosed Hypertension Patients	
Widya Warastuti (Author)	556-561
□ PDF	
Molecular Docking Analysis of Ficus religiosa Active Compound with Anti-Inflammatory Activity by Target Tumour Necrosis Factor Alpha and Vascular Endothelial Growth Factor Receptor in Diabetic Wound Healin	_
Yuyun Yueniwati, Mokhammad Fahmi Rizki Syaban, Nabila Erina Erwan, Gumilar Fardhani Ami Putra, Agung Dwi Krisnayana (Author)	1031-1036
□ PDF	
The Effectiveness of Binahong (Anredera cordifolia (Ten.) Steenis) Extract in Promoting Fertility in Male W Rats after Exposure to Cigarette Smoke	istar
Achmad Ramadhan (Author)	123-128
□ PDF	
On the Issue of Creating a Cream of Reparative Action for the Treatment and Prevention of Cheilitis	
Alexander Kotenko, Tatyana Yarnykh, Natalya Zhivora, Olga Rukhmakova, Vladimir Kovalev, Marina Buryak (Author)	308-312
□ PDF	
The Effect of Ant Plant on the Increase of CD4 Count in PLHIV in the Papua Province	
Arwam Hermanus Markus Zeth, Nouvy Helda Warouw, Paula Krisanty (Author)	1231-1239
□ PDF	
Photoprotective and Inhibitory Activity of Tyrosinase in Extract and Fractions of Terminalia catappa L.	
Maulita Indrisari, Sartini Sartini, Upik A. Miskad, Khairuddin Djawad, Karlina Amir Tahir, Nurkhairi Nurkhairi, Lukman Muslimin (Author)	263-270
₽DF	
Brine Shrimp (Artemia salina Leach.) Lethality Test of Ethanolic Extract from Green Betel (Piper betle Linn Red Betel (Piper crocatum Ruiz and Pav.) through the Soxhletation Method for Cytotoxicity Test	.) and
Nerdy Nerdy, Puji Lestari, Jon Piter Sinaga, Selamat Ginting, Nilsya Febrika Zebua, Vriezka Mierza, Tedy Kurniawan Bakri (Author)	407-412
PDF	
Antihyperglycemic, Endothelial protection and Toxicity study of Basil Leaves Extract on Diabetic Rats	
Sry Suryani Widjaja, Rusdiana Rusdiana (Author); Maya Savira; Rina Amelia (Author)	589-594
□ PDF	

Tyrosinase Enzymes Activities and Sun Protection Factor of Ethanol Extract, Water Fraction, and n-Butanol Fraction of Chromolaena odorata L. Leaves

Karlina Amir Tahir, Upik A. Miskad, Khairuddin Djawad, Sartini Sartini, Natsir M. Djide, Maulita Indrisari, Khaerani Khaerani, Syamsuri Syakri, Anshari Masri, Ahmad Lalo (Author)

493-498

☑ PDF

Silver Nanoparticle of Acalypha indica Linn. Leaf As Bio-larvicide against Anopheles sp. Larvae

Yos Banne, Olfie Sahelangi, Steven Soenjono, Elisabeth Natalia Barung, Selfie Ulaen, Rivolta G. M. Walalangi, Zulfiayu Sapiun (Author)

760-765

☑ PDF

Characterization and Anti-aging Tests of Peel-Off Gel Masks Made from Ethanolic Extract of Yarrow (Achillea millefolium)

Syamsuri Syakri, Isriany Ismail, Nurul Muamanah Amal, Nur Asma Masjidi, Karlina Amir Tahir (Author)

1156-1161

PDF

Antioxidant and Hypolipidemic Effects of Ipomoea batatas L and Pandanus conoideus Lam Combination on Rats Fed with High Cholesterol Diet

A. A. Ngurah Subawa, I Wayan Putu Sutirta Yasa, I Made Jawi, Agung Nova Mahendra (Author)

473-476

☑ PDF

Dewan Redaksi

Open Access Macedonian Journal of Medical Sciences

About ▼ Announcements Metrics ▼ Current ▼ Archives EU ▼ Scopus ▼ NCBI ▼

Archives EU → Scopus → NCBI → CABI → Indexation →

n 🕶 🔾 Search

Register

Login

No Publication Fee: ▼ Membership ▼

Home / Editorial Team

Editorial Team

Journal Manager

Dr. Eli Djulejic, Open Access Macedonian Journal of Medical Sciences, Belgrade, Serbia

Editor-in-Chief

Prof. Dr. Mirko Zhivko Spiroski, Scientific Foundation SPIROSKI, Rajko Zhinzifov No 48, 1000 Skopje, Republic of Macedonia

Section Editors (Deputy Editors-in-Chief)

Dimitrov Borislav D, MD. Academic Unit of Primary Care and Population Sciences Faculty of Medicine University of Southampton South Academic Block (Level C) Southampton General Hospital Southampton SO166YD England, UK

 ${\it Mukaetova-Ladinska~Elizabeta,~MD,~PhD,~MRCP sych.~Old~Age~Psychiatry,~Newcastle~University,~United~Kingdom~Age~Psychiatry,~Newcastle~University,~United~Kingdom~Age~Psychiatry,~Newcastle~University,~United~Kingdom~Age~Psychiatry,~Newcastle~University,~United~Kingdom~Age~Psychiatry,~Newcastle~University,~United~Kingdom~Age~Psychiatry,~Newcastle~University,~United~Kingdom~Age~Psychiatry,~Newcastle~University,~United~Kingdom~Age~Psychiatry,~Newcastle~University,~United~Kingdom~Age~Psychiatry,~Newcastle~University,~United~Kingdom~Age~Psychiatry,~Newcastle~University,~United~Kingdom~Age~Psychiatry,~Newcastle~University,~United~Kingdom~Age~Psychiatry,~Newcastle~University,~United~Kingdom~Age~Psychiatry,~Newcastle~University,~United~Kingdom~Age~Psychiatry,~Newcastle~University,~United~Kingdom~Age~Psychiatry,~Newcastle~University,~United~Kingdom~Age~Psychiatry,~Newcastle~University,~Newcastle~$

Branislav Filipović, MD, PhD. University of Belgrade, Faculty of Medicine, Institute of Anatomy "Niko Miljanicâ€□ Address: 4/2 Dr Subotica Starijeg, Belgrade, Serbia

Spiroski Igor, MD. University Clinic of Cardiology, Faculty of Medicine, Ss Cyril and Methodius University of Skopje, Skopje, Republic of Macedonia

Stoleski Sasho, MD, PhD. Institute for Occupational Health of Republic of Macedonia - Skopje, WHO Collaborating Center, GA2LEN Collaborating Center, II Makedonska brigada 43, 1000 Skopje, Republic of Macedonia

Hristomanova-Mitkovska Slavica, MD, MSc. Institute for Human Genetics of the Faculty of Medicine in Göttingen, Germany

Bogoeva Ksenija, MD, PhD. PHO Prof Bogoev, Skopje, Republic of Macedonia

Stojanovski, Sinisa, MD, PhD.

Information

For Readers

For Authors

For Librarians

Open Journal Systems

Make a Submission

Browse

Categories

- A Basic Sciences
- B Clinical Sciences
- C Case Reports
- D Dental Sciences
- E Public Health
- F Review Articles
- G Nursing
- T Thematic Issues
- T1 "Coronavirus Disease (COVID-19)"
- T2 "Public Health and Nutrition Sciences in the Current Millennial Era"
- T3 "Neuroscience, Neurology, Psychiatry and General Medicine"
- T4 "Contribution of Nurses on Sustainable Development Goals
- T5 "Re-Advancing Nursing Practice, Education and Research in the Post Covid"

Layout Editor and Electronic Publishing

MSc, Eng Ivo Spiroski, ID Design 2012, Skopje, Republic of Macedonia

Copyeditor

Sinjore, India

Evidence Based Medicine

Prof. Dr. Katarina Stavric, Children Hospital Skopje, Macedonia, Vodnjanska 17, University "Ss Cyril and Methodius", Skopje, Republic of Macedonia

Editorial Board

DDS, MS, PhD, Associate Professor Nikola Angelov, Director of the Pre-Doctoral Periodontics Clinic, Loma Linda University School of Dentistry, Department of Periodontics. Loma Linda, CA, 92350, United States

Assist. Prof. Dr. Ramush Bejiqi, University Clinical Centre of Kosovo, Paediatric Clinic, Albania

Prof. Semra ÄŒavaljuga, Department of Epidemiology and Biostatistics, Faculty of Medicine, Sarajevo, Bosnia and Herzegovina

MD Pei-Yi Chu, Diagnostic and research pathologist, Department of Surgical Pathology, Changhua Christian Hospital, Taiwan. Address: 135 Nan-Shiao Street, Changhua 500-06,, Taiwan, Province of China

MD, PhD Ivo Donkov, Staff Urologist, Lincoln County Hospital, United Kingdom

MD, PhD Andrew J. Dwork, Departments of Pathology and Cell Biology and Psychiatry, College of Physicians and Surgeons of Columbia University; Division of Molecular Imaging and Neuropathology, New York State Psychiatric Institute, Unit 62, 722 West 168th Street, New York, NY 10032, United States

Adriana Galan, Department of Health Programmes and Health Promotion, Institute of Public Health, Bucharest, Romania

Prof. Tania Santos Giani, Estacio de Sa University, in Health Sciences, Brazil

PhD Iva Ivanovska, Harvard Medical School, Department of Genetics, 77 Avenue Louis Pasteur, NRB room 239, Boston, MA 02115. United States

MD, PhD Jerzy JabÅ, ecki, Associate Professor, Division of General Surgery St. Jadwiga of Silesia Hospital, Trzebnica; Head, Subdepartment of Hand Surgery an Replantation St Jadwiga of Silesia Hospital, Trzebnica; Professor, Department of Public Health, State Higher Professional Medical School, Opole, Poland. 55-100 Trzebnica, ul. Prusicka 53, Poland

MD Mehrdad Jalalian Hosseini, Khorasan-e Razavi Blood Center, Mashhad, Iran, Islamic Republic of

PhD Radka Kaneva, Department of Medical Chemistry and Biochemistry, Medical University - Sofia, Bulgaria

Prof. Dr. Kostandina Leonida Korneti-Pekevska, Ss Cyril and Methodius University of Skopje, Faculty of Medicine, Skopje, Republic of Macedonia

MD, PhD Branko Malenica, Department of Immunology, Clinical Hospital Center Zagreb, Zagreb University School of Medicine, Zagreb, Croatia

Prof. Dr. Elida Mitevska, Institute of Histology and Embriology, Faculty of Medicine, Ss Cyril and Methodius University of Skopje, Skopje, Republic of Macedonia

T6 - "The Chalenges and Opportunities for Nurses in The New Era Adaptation"

T7 - "APHNI: Health Improvement Strategies Post Pandemic Covid-19"

T8 -"Pharmacy collaboration in achieving health resilience"



Part of the PKP Publishing Services Network

MD, PhD Jerzy Jabå,ecki, Associate Professor, Division of General Surgery St. Jadwiga of Silesia Hospital, Trzebnica; Head, Subdepartment of Hand Surgery an Replantation St Jadwiga of Silesia Hospital, Trzebnica; Professor, Department of Public Health, State Higher Professional Medical School, Opole, Poland. 55-100 Trzebnica, ul. Prusicka 53, Poland

MD Mehrdad Jalalian Hosseini, Khorasan-e Razavi Blood Center, Mashhad, Iran, Islamic Republic of

PhD Radka Kaneva, Department of Medical Chemistry and Biochemistry, Medical University - Sofia, Bulgaria

Prof. Dr. Kostandina Leonida Korneti-Pekevska, Ss Cyril and Methodius University of Skopje, Faculty of Medicine, Skopje, Republic of Macedonia

MD, PhD Branko Malenica, Department of Immunology, Clinical Hospital Center Zagreb, Zagreb University School of Medicine, Zagreb, Croatia

Prof. Dr. Elida Mitevska, Institute of Histology and Embriology, Faculty of Medicine, Ss Cyril and Methodius University of Skopje, Skopje, Republic of Macedonia

MD, PhD Marija Mostarica-Stojković, Institute of Microbiology and Immunology, University of Belgrade School of Medicine, Belgrade, Serbia

PhD Vesna Nikolova-Krstevski, Harvard Institutes of Medicine, HIM-201, 4 Blackfan Circle, Boston, MA, 02134, United States

Prof. Dr. Nikola Panovski, Institute of Microbiology and Parasitology, Faculty of Medicine, Skopje, Republic of Macedonia

MD, BIDMC Iva Petkovska, Beth Israel Deaconess Medical CenterRadiology W CC - 3 330 Brookline Ave. Boston, MA 02215, United States

Prof. Dr. Gordana Petrusevska, Institute of Pathologyy, Medical Faculty, University of "Ss. Cyril and Methodiusâ€□ â€″ Skopje, Republic of Macedonia

Prof. Enver Roshi, Dean of Faculty of Public Health, Medical University of Tirane, Chief of Epidemiological Observatory, National Institute of Public Heath. Address: Rruga e Dibres, Str. 371, Tirana, Albania

MD, PhD Gorazd B. Rosoklija, Professor at Columbia University and member of the Macedonian Academy of Sciences and Arts. United States

Prof. Dr. Aleksandar Sikole, University Clinic for Nephrology, Faculty of Medicine, Ss Cyril and Methodius University of Skopje, Skopje, Republic of Macedonia

MD, PhD Rumen Stefanov, Information Centre for Rare Diseases and Orphan Drugs (ICRDOD), Bulgaria; Department of Social Medicine, Medical University of Plovdiv, Bulgaria

Prof. Dr. Vesna Velikj Stefanovska, Department of Epidemiology and Biostatistics with Medical Informatics, Medical Faculty, UKIM, Skopje, Republic of Macedonia

MD, MBA Milenko Tanasijevic, Director, Clinical Laboratories Division and Clinical Program Development, Pathology Department, Brigham and Women's Hospital, Dana Farber Cancer Institute, Associate Professor of Pathology, Harvard Medical School, United States

MD, FRCPC Kiril Trpkov, Associate Professor, University of Calgary, Department of Pathology and Laboratory Medicine, Calgary Laboratory Services. 7007 14 st, Calgary SW, Canada

MD, PhD Igor Tulevski, Department of Cardiology, Academic Medical Center, Amsterdam, 1100 DD, T 020 707 2930; F 020 707 2931, Netherlands

Past Members of Editorial Team

Prof. Dr Doncho Donev, Institute of Social Medicine, Medical Faculty, Ss Cyril and Methodius University of Skopje, Skopje, Republic of Macedonia

Prof. Dr Olivera Stojceva Taneva, University Clinic of Nephrology, Republic of Macedonia

Prof. Dr Petar Miloshevski, from 2008-2014, Institute of Preclinical and Clinical Pharmacology with Toxicology, Faculty of Medicine, Skopje, Republic of Macedonia

Prof. Dr Sonja Topuzovska, Institute of Medical and Experimental Biochemistry, Faculty of Medicine, Skopje, Republic of Macedonia

Prof. Dr. Aleksandar Dimovski, Institute of Pharmaceutical Chemistry, Faculty of Pharmacy, University "Ss Kiril and Metodij", Skopje, Republic of Macedonia

PhD Mirko Trajkovski, ETH Zurich, Wolfgang-Pauli-Str. 16/HPT D57, 8093 Zurich-CH, Switzerland

PhD Zoran Zdravkovski, Institute of Chemistry, Faculty of Natural Sciences and Mathematics, Ss Cyril and Methodius University of Skopje, Skopje, Republic of Macedonia

Published by



About us

About the Journal
Editorial Team
Author Fees
Online Payments
Bibliographic Information
Journal History

Information

For readers
For authors
For librarians
For Reviewers
Transparency

Contact

Publisher
Co publisher
Registration
Contact
Privacy statement

Platform & workflow by OJS / PKP

Scientific Foundation SPIROSKI, Skopje, Republic of Macedonia Open Access Macedonian Journal of Medical Sciences. 2021 Sep 17; 9(A):760-765 https://doi.org/10.3889/oamjms.2021.6628 elSSN: 1857-9655

Category: A - Basic Sciences

Section: Pharmacology





Silver Nanoparticle of Acalypha indica Linn. Leaf As Bio-larvicide against Anopheles sp. Larvae

Yos Banne¹*10, Olfie Sahelangi²10, Steven Soenjono³10, Elisabeth Natalia Barung¹10, Selfie Ulaen¹10, Rivolta G. M. Walalangi² D, Zulfiayu Sapiun⁴

¹Department of Pharmacy, Politeknik Kesehatan Kementerian Kesehatan, Manado, Indonesia: ²Department of Nutrition, Politeknik Kesehatan Kementerian Kesehatan, Manado, Indonesia; ³Department of Environmental Health, Politeknik Kesehatan Kementerian Kesehatan, Manado, Indonesia; ⁴Department of Pharmacy, Politeknik Kesehatan Kementerian Kesehatan, Gorontalo, Indonesia

Abstract

BACKGROUND: Acalypha indica Linn, has been used as traditional medicine, it contains flavonoids, alkaloids, tannins, saponins, steroids, triterpenoids, and essential oils.

AIM: This study aimed to determine the bio-larvicide effects of A. indica Linn. leaf stew and the silver nanoparticles against Anopheles sp. larvae.

METHODS: The fresh leaves of A. indica Linn, extracted using distilled water at 100°C for 30 min. The silver nanoparticles were made by mixing a solution of silver nitrate with the stew, which acts as a reducing agent. The resulting silver nanoparticles were characterized by particle size analyzer and UV-vis spectrophotometer. The bio-larvicide effects against Anopheles sp. larvae performed using a completely randomized design. There were eight groups consisted of ten larvae and three replications. Treatment groups of stew and silver nanoparticle for concentrations 0.05%, 0.5%, and 5%, respectively. The negative control group was distilled water and the positive control group was the 0.01% abate solution. Assessment of larvicide activity was carried out every hour for 6 h and continued if there were larvae that live up to 24 h. The LC₅₀ value was calculated based on Probit

RESULTS: The results showed that the A. indica Linn. leaf stew can be made into silver nanoparticles preparations, optimal results were obtained from a mixture of 1% stew and 3 mM AgNO₃. The result of bio-larvicides effect test against Anopheles sp. larvae showed that the LC₅₀ value of the A. indica Linn. leaf stew was 727,3 ppm and the LC₅₀ value of silver nanoparticles was 3.366 ppm.

CONCLUSION: It can be concluded that A. indica Linn. is a promising larvicidal plant and can be made into silver nanoparticle preparations

Edited by: Mirko Spiroski Citation: Banne Y, Sahelangi O, Soenjono S, Barung EN, Ulaen S, Walalangi RGM, Sapiun Z. Silver Nanoparticle of Acalypha indica Linn. Leaf As Bio-larvicide against Anopheles sp. Larvae, Open Access Maced J Med Sci. 2021 Sep 17; 9(A):760-765. https://doi.org/10.3889/oamjms.2021.6628 Keywords: Acalypha indica Linn. leaf; Stew; Silver nanoparticles; Bio-larvicide; Anopheles sp. *Correspondence: Yos Banne, Department of Pharmacy Politeknik Kesehatan Kementerian Kesehatan

Politeknik Kesehatan Kementerian Kesehatan, Manado, Jl. R.W. Monginsidi Malalayang II, Manado, Indonesia. E-mail: yosbanne.250108@gmail.com Received: 14-Jun-2021 Revised: 15-Aug-2021 Accepted: 07-Sep-2021 Copyright: © 2021 Yos Banne, Olfie Sahelangi,

Steven Soenjono, Elisabeth Natalia Barung, Selfie Ulaen, Rivolta G. M. Walalangi, Zulfiayu Sapiun Funding: This study was supported by Politeknik Kesehatan Kementerian Kesehatan Manado Competing Interests: The authors have declared that no competing interests exist

Open Access: This is an open-access article distributed under the terms of the Creative Commons Attribution NonCommercial 4.0 International License (CC BY-NC 4.0)

Introduction

Malaria disease is caused by the Plasmodium parasite, which is spread by female Anopheles mosquitoes. According to the WHO report, globally, there were 229 million cases in 2019 with a mortality rate 10 in 100.000 population at risk [1]. Handling malaria can be done with prevention and treatment. Prevention can be done using insect repellent lotion, removing breeding places and standing water around the house, and killing mosquito larvae by fogging or using larvicides [2]. Chemical substances and natural materials can be used as larvicides. Research on the larvicidal effect has been carried out on several plants for example Acalypha indica Linn. [3], [4], Solanum xanthocarpum, Euphorbia tirucalli, Momordica charantia, Eucalyptus globules, Citrullus colocynthis, A. indica, Annona squamosa, and Solanum nigrum [5].

A. indica Linn. is a type of weed wild plants that are often found on the roadside, untreated grass fields even as a nuisance on agricultural land [6]. According to Hayati et al., the components contained in this plant are β -sitosterol and daucosterol, saponins, tannins, flavonoids, and essential oils [7]. The community has been used this plant as traditional medicine and the results of the researches prove its efficacy, among others, as antidiabetic, antiinflammatory, antioxidant, anti-poison, anticancer, hepatoprotective, antibacterial, wound healing, anthelmintic, repellent, and larvicide [3], [8], [9], [10]. Research conducted by Pratiwi et al. (2015) shows that the ethyl acetate extract of A. indica Linn. has a larvicidal effect on Aedes aegypti mosquito larvae with an LC_{50} of 72.444 ppm [10].

Pharmacological effects possessed by natural substances are produced by the content of secondary metabolites. Several methods can be used to find the content of natural ingredients, generally distinguished by the hot method and cold method. The choice of method is based on the physical and chemical properties of the secondary metabolites to be explored. The decoction is one of the simple extraction methods using water as a

solvent; the material to be extracted must be resistant to heating [12].

Several studies that use natural ingredients try to compare the pharmacological effects produced between plant extracts and plant nanoparticle preparations. The results obtained indicate that an increase in the effect obtained from the preparation of nanoparticles, due to the small particle size of the drug which allows the drug to be absorbed and penetrate quickly into cells and provide the therapeutic effect. Silver nanoparticles are one of the nanoparticle preparations that have been developed. Nano-sized silver metal particles have antimicrobial properties but are environmentally friendly and safe for humans [13], [14]. The green synthesis method uses plant extracts that act as reducing agents which react with metal ions to produce silver nanoparticles. Plant parts can be used in the form of leaves, fruit, or seeds [14]. This study aimed to make the A. indica Linn. leaf stew and silver nanoparticle preparations, and determine the bio-larvicide effect against Anopheles sp. larvae.

Materials and Methods

Materials

A. indica Linn. leaves were collected in June from Malalayang district at Manado City. AgNO₃ purchased from Merck Company. Polyvinyl alcohol (PVA) purchased from Sigma-Aldrich. Abate purchased from BASF Indonesia. Test animals were Anopheles sp. third instar larvae purchased from Department of Environmental Health, Politeknik Kesehatan Kementerian Kesehatan Manado, Indonesia.

Methods

Stew and silver nanoparticles preparation

- 1. Stew preparation: *A. indica* Linn. leaves taken in the morning and washed with running water, and then the leaves were air-dried. Furthermore, the leaves were coarsely powdered using a grinder. The leaves powder weighed 20 g and placed in Erlenmeyer containing 200 mL of distilled water, then simmer for 30 min. Then the solution was cooled to room temperature then filtered with Whatman filter paper no. 1, and the stew was stored in a cold storage cabinet at 4°C. For the nanoparticle preparation, the stew was diluted with aquadest to obtain a concentration of 0.5, 1, and 5% [15].
- Silver nanoparticle preparation: Optimization of the nanoparticles preparation carried out by

adding 200, 400, 600, 800, 1000, 1200, 1400, 1600, 1800, and 2000 µL of 1% *A. indica* Linn. leaf stew to 1 mL of 1 mM AgNO₃ solution. Also made preparations using variations in the concentration of AgNO₃ 1–5 mM added with the 0.5 and 5% *A. indica* Linn. leaf stew. As a stabilizer 0.5% PVA solution was added for every 2 mL AgNO₃. The mixture was allowed to stand in a dark container at cool temperatures for 24 h. The formation of silver nanoparticles was characterized by a change in color of the solution from colorless to slightly turbid brown with wavelengths of 430–480 nm and particle size of 100 nm ≥[15]. The silver nanoparticle preparation then processed using a freeze dryer.

- 3. Characterization of silver nanoparticle:
- UV-Vis spectrophotometer
 UV-Vis spectrophotometer was standardized
 using blank. The extract solution was put in a cuvette and then measured at a wavelength of 350–500 nm [14].
- b. Particle size and morphology analysis
 The particle size of the extract preparation measured using a Particle Size Analyzer (PSA) [14], [16]. The formulation that has the smallest particle size with the appropriate wavelength was then used in the bio-larvicide test.

Secondary metabolites test

The test of the secondary metabolites content in the stew was carried out qualitatively using chemical reagents to determine alkaloid, tannin, terpenoid, steroid, flavonoid, and saponin.

Bio-larvicide test

A total of 10 Anopheles sp. third instar larvae transferred from the container into a vial containing the A. indica Linn. stew with a concentration of 0.05%, 0.5%, and 5%, control (+) by giving an abate concentration of 0.01%, and control (-) by giving distilled water. Mosquito larvae activity of Anopheles sp. was observed for 24 h. Observation of dead larvae was carried out at the hours 1, 2, 3, 4, 5, and 6. Observation of the flow of life was continued if there were larvae that lived until 24 h after treatment. Calculation of time begins after the insertion of larvae into glass beakers. Observation of the life cycle, that is, the test larvae were given extracts able to survive for a certain period of time but could not reach the next stage. Larvae are considered dead if there are no more signs of life, for example, do not move anymore even though stimulated by the movement of water and touched with a stick [11].

The data obtained were made in the form of tables and graphs and LC_{50} values were calculated

A - Basic Sciences Pharmacology

based on Probit analysis.

secondary metabolites produce pharmacological effects.

Results

Stew and silver nanoparticles preparation

From 110 samples with varying concentrations of *A. indica* Linn. stew and AgNO₃, there were six samples that met the criteria of wavelength 430–480 nm and particle size on the nanometer scale, as shown in Table 1 and Figure 1.

Table 1: Optimization results of silver nanoparticles of *Acalypha indica* Linn. leaves

Sample	Mix of	λ (nm)	Particle size (nm)			
code			D1	D2	D3	Mean
B1	AILS 1% - AgNO ₃ 3 mM	460, 449, 441	106,7	100,6	104,9	104,1
B2	AILS 1% - AgNO ₃ 3 mM	446	88,2	96,8	108,2	97,7
B3	AILS 1% - AgNO ₃ 3 mM	451	91,8	82,5	91,8	88,7
B4	AILS 1% - AgNO 3 mM	442	100,8	101,5	99,6	100,6
F1	AILS 0,5% - AgNO ₃ 5 mM	431	130,0	128,1	129,6	129,2
G3	AILS 1% - AgNO ₃ 5 mM	468	93,1	92,2	91,8	92,4

AILS: Acalypha indica Linn. leaves stew

Based on these results, those who met the criteria as silver nanoparticles *A. indica* Linn. leaves was B3 samples because they have the smallest particle size.



Figure 1: Acalypha indica Linn. leaves stew (a), silver nanoparticles of A. indica Linn. leaves stew (b)

Secondary metabolites test

The test conducted using chemical reagents to determine alkaloid, tannin, terpenoid, steroid, flavonoid, and saponin in the stew. The result showed that all the secondary metabolites tested were contained in the stew (Table 2). The content of these

Table 2: Secondary metabolites in the stew of Acalypha indica Linn. leaves

Secondary metabolites test	Result
Alkaloid	+
Saponin	+
Tannin	+
Flavonoid	+
Steroid	+
Terpenoid	+
+: Contains the compound, -: Non contains the compound.	-

Bio-larvicide test of stew and silver nanoparticles of A. indica Linn. leaf stew

The results of the bio-larvicides test of *A. indica* Linn. leaf stew and silver nanoparticles of *A. indica* Linn. leaf stew against the larvae of *Anopheles sp.* Are presented in Table 3.

Table 3: Bio-larvicides effects of stew and silver nanoparticles of *Acalypha indica* Linn. leaf stew in the larvae of *Anopheles* sp.

Sample	Mortality percentage of Anopheles sp. mosquito larvae			
	N1	N2	N3	Mean
Positive control	100	100	100	100
Negative control	0	0	0	0
Stew 0,05%	60	50	50	53,33
Stew 0,5%	80	80	70	73,33
Stew 5%	100	100	100	100
Silver Nanoparticle 0.05%	90	90	90	90
Silver Nanoparticle 0.5%	100	100	100	100
Silver Nanoparticle 5%	100	100	100	100

Based on these data, the LC_{50} value is calculated based on the Probit analysis as follows in Table 4.

Table 4: Probit analysis data for LC₅₀ calculations

Sample	Concentration (%)	Log	% Mortality	Probit value
Stew	0.05	-1.30103	53.3	5.08
Stew	0.5	-0.30103	73.3	5.61
Stew	5	0.69897	100	8.09
Nanoparticle	0.05	-1.30103	90	6.48
Nanoparticle	0.5	-0.30103	100	8.09
Nanoparticle	5	0.69897	100	8.09

From the regression equation, the LC_{50} value for *A. indica* Linn. leaf stew is 0.07273%, if the value is converted to the weight of the *A. indica* Linn. leaf it will be equivalent to 10 fresh leaves/100 mL. While, the LC_{50} value for nanoparticles of *A. indica* Linn. stew leaf is 0.0003366%.

Discussion

Samples of *A. indica* Linn. leaves used in dry form. The drying process by air-dried to prevent damage to the content in the leaves due to the sunlight. Stew is one of the simple methods of extracting heat using water as a solvent [11]. The choice of method and solvent is based on the physical and chemical properties of the secondary metabolites to be explored. Polar materials such as flavonoids, saponins, and tannins are extracted with polar solvents such as water and n-butanol [17]. Most flavonoid content is found in the leaves of this plant [18]. The stew was obtained in the form of dark brown liquid with a specific aroma like *A. indica* Linn.

To determine the right formulation of silver nanoparticles, the optimization of the preparations

was done using various comparisons of several concentrations of stew and ${\rm AgNO_3}$ solution. The formation of silver nanoparticles is characterized by a brownish yellow color as seen in Figure 1. To ensure, the measurement of the wavelength is compared with the stew of the *A. indica* Linn. leaf. Silver nanoparticles have a wavelength of 430–480 nm, while the stew has a wavelength of 300–380 nm. The particle diameter measurement is done using the PSA. The sample chosen for testing the effects of bio-larvicides was those that had the smallest particle size. The small particle size allows the drug to absorb and penetrate rapidly into cells and provide a therapeutic effect.

The secondary metabolite test conducted using chemical reagents. The result showed that alkaloid, tannin, terpenoid, steroid, flavonoid, and saponin found in the stew. The previous study results showed that this plant contains β -sitosterol and daucosterol, saponins, tannins, flavonoids, and essential oils [7]. This plant has been used in traditional medicine as antidiabetic, anti-inflammatory, antioxidant, anticancer, hepatoprotective, antibacterial, wound healing, anthelmintic, and larvicide [8], [9].

The bio-larvicide test uses Anopheles sp. third instar larvae and observations are carried out every hour for 6 h and if there are still mosquitoes that do not die then the observations are carried out for up to 24 h to observe the life path. The third instar larvae were chosen for the test because the size is larger than the first and the second instar larvae so that the observation becomes easier. Besides that, the third instar larvae are more resistant to current mechanical factors [10]. The results showed that the greater the stew concentration, the greater the percentage of mosquito larvae mortality and the larvae that did not die while the experience period would not grow to the next phase. Komalamisra et al. [19] considered larvicidal products exerting LC_{50} <50 mg/L (active), 50 mg/L <LC $_{50}$ <100 mg/L (moderately active), $100 \text{ mg/L} < LC_{50} < 750 \text{ mg/L} (effective), and LC_{50} > 750 \text{ mg/L}$ (inactive). Kiran et al. [20] considered compounds with LC₅₀ <100 mg/L as exhibiting a significant larvicidal effect.

Phytochemicals such as alkaloids, phenols, and triterpenoids alone or in combination, contribute to acute toxicity toward various arthropod species [21]. The death of mosquito larvae in this research is presumed due to the secondary metabolites content in A. indica Linn. leaves, that is, alkaloid, triterpenoid, flavonoid, tannin, saponin, and steroid, Alkaloid compounds work by disrupting the system nerves (neuromuscular toxic), inhibiting the feeding power of the larva, and acts as a stomach poison. The mechanism of action of the alkaloids, namely by inhibiting the action of the enzyme functioning acetylcholinesterase hydrolyzes acetylcholine. In a state of acetylcholine normally conducts function nerve implants, after which it is immediately experienced hydrolysis with the help of acetylcholinesterase into choline and acetic acid. With the help of this acetylcholinesterase enzyme occurs build-up of acetylcholine to be causes disturbance and damage in the nervous system [22]. Alkaloids also act as an antifeedant and stomach poisoning [23]. Flavonoids act as an insecticide by attacking the nerves in some vital organs of insects, thus arising a weakening of the nerves such as the inhalation resulting in the larvae unable to breathe and finally died. Flavonoids work as a respiratory inhibitor and are considered to disturb the energy metabolism in the mitochondria by inhibiting the conveying system electrons [24]. Triterpenoids are the active components of the limonoid compounds that are toxic used as an insecticide as potential as an antifeedant against insects, growth regulators, and substances toxic to rice lice, larvicides, antimicrobial, insect repellent, and reproductive inhibitor [25]. Steroids have a chemical structure similar to triterpenoids. Steroids are hormone growth that affects the molting of the larvae. Steroids will cause chitin cell walls in the body larvae to thicken; resulting in larval growth will be disturbed and cause death in larvae [26]. Saponins cause a decrease in the activity of digestive enzymes and the absorption of food in insects [27]. In addition, saponins also damage the cuticle membrane of larvae and cause the death of larvae [28]. The tannin compounds contained to act as digestive toxins and interfere with water absorption in larvae so that they can cause death [27], [29].

The LC_{50} value shows that nanoparticle preparation has much stronger bio-larvicide effects than the stew. This is possible because of its small size making it easier to penetrate cells, as well as the presence of nanosized silver metal content that has antimicrobial properties but is environmentally friendly and safe for humans [13]. The results obtained in this study indicate the potential for the development of stew and silver nanoparticles of the A. indica Linn. leaves to be used as bio-larvicides.

Conclusions

 $A.\ indica$ Linn. leaves can be extracted with a simple method of stew and the results of the stew can be made into silver nanoparticle preparations. $A.\ indica$ Linn. is a promising larvicidal plant in which the LC $_{50}$ value of the leaf stew and its silver nanoparticles were 727.3 ppm and 3.366 ppm, respectively. It is recommended to conduct similar research on mosquito larvae that cause other diseases such as $A.\ aegypti$ and Culex sp. as well as conducting further research on the stability of silver nanoparticle preparations.

Acknowledgment

The authors are thankful to the director of Politeknik Kesehatan Kementerian Kesehatan Manado

A - Basic Sciences Pharmacology

for funding support and providing facilities for the research; to the management of the Nanopharmacy Research Center Laboratory of the Department of Pharmacy, Universitas Islam Indonesia for providing facilities for nanoparticle preparation.

Credit Author Statement

YB: Conceptualization, Methodology, Investigation, Writing-Original draft, Writing- Reviewing and Editing; OS: Visualization, Validation; EB: Formal analysis, Supervision; SS: Resources, Methodology; RW: Resources, Conceptualization; SU: Project administration, Data curation; ZS: Writing- Reviewing and Editing, Data curation.

References

- World Health Organization. World Malaria Report 2020, Years of Global Progress and Challenges. Geneva: World Health Organization; 2021. Available from: https://www.who.int. [Last accessed on 2021 Mar 06].
- Tizifa TA, Kabaghe AN, McCann RS, Berg HV, Vugt MV, Phiri KS. Prevention efforts for malaria. Curr Trop Med Rep. 2018;5(1):41-50. https://doi.org/10.1007/s40475-018-0133-y PMid:29629252
- Teklani PW, Perera BG. Mosquito repellent and larvicidal activities of Acalypha indica leaf extracts. Int J Pharm Pharmacol. 2017;1(2):107.
- Karthik S, Suriyaprabha R, Vinoth M, Srither SR, Manivasakan P, Rajendran V, et al. Larvicidal, super hydrophobic and antibacterial properties of herbal nanoparticles from Acalypha indica for biomedical applications. RSC Adv. 2017;7:41763-70. https://doi.org/10.1039/c7ra05697d
- Gosh A, Chowdhury N, Chandra G. Plant extracts as potential mosquito larvicides. Indian J Med Res. 2012;135(5):581-98.
 PMid:22771587
- Pambudi A, Syaefuddin, Noriko N, Swandari R, Azura PR. Identification of flavonoid bioactives from *Acalypha indica* L. J Al-Azhar Indones Seri Sains Teknol. 2014;2(3):178-87. https://doi.org/10.36722/sst.v2i3.139
- Hayati EK, Jannah A, Identification of compounds and antimalarial activity in vivo ethyl acetate extract of the plant Acalypha indica L. Molekul. 2012;7(1):20-32. https://doi. org/10.20884/1.jm.2012.7.1.103
- Nahrstedt A, Hungeling M, Petereit F. Flavonoids from Acalypha indica. Fitoterapia. 2006;77(6):484-6. https://doi. org/10.1016/j.fitote.2006.04.007
 PMid:16828241
- Duke JA. List of chemicals of Acalypa australis L. In: Phytochemical and Ethnobotanical Databases; 2009. Available from: http://www.sun.arsgrin.gov:8080/npgspub/xsql/plantdisp.xsql?taxon=406. [Last accessed on 2021 Mar 06].

 Zahidin NS, Saidin S, Zulkifli RM, Muhamad II, Ya'akob H, Nur H. A review of *Acalypha indica* L. (Euphorbiaceae) as traditional medicinal plant and its therapeutic potential. J Ethnopharmacol. 2017;207:146-73. https://doi.org/10.1016/j. jep.2017.06.019

PMid:28647509

- Pratiwi D, Prahastiwi EA, Savitri M. Larvicidal activity test of Acalypha indica Linn herb ethyl acetate extract. against Aedes aegypti mosquito larvae. Farmagazine 2015;2(1):16-23. https://doi.org/10.20884/1.jm.2012.7.1.103
- Depkes RI. Direktorat Jenderal Pengawasan Obat dan Makanan. 4th ed. Indonesia: Farmakope Indonesia; 1995. p. 551, 713.
- Tolaymat T, El Badawy A, Genaidy A, Scheckel K, Luxton T, Suidan M. An evidence-based environmental perspective of manufactured silver nanoparticle in syntheses and applications: A systematic review and critical appraisal of peer-reviewed scientific papers. Sci Total Environ. 2010;408(5):999-1006. https://doi.org/10.1016/j.scitotenv.2009.11.003

PMid:19945151

- Nalawati AN. Sintesis Nanopartikel Perak (NPAg) Dengan Metode Yang Ramah Lingkungan Dan Kajian Aktivitasnya Dalam Menghambat Pertumbuhan Bakteri Gram Positif dan Bakteri Gram Negatif [Tesis]. Bogor: Institut Pertanian Bogor; 2015. https://doi.org/10.6066/jtip.2017.28.1.27
- Ahmed S, Ullah S, Ahmad M, Swami BL, Ikram S. Green synthesis of silver nanoparticles using *Azadirachta indica* aqueous leaf extract. J Radiat Res Appl Sci. 2016;9(1):1-7. https://doi.org/10.1016/j.jrras.2015.06.006
- Abdel-Azis MS, Shahen MS, El-Nekeety A, Abdel-Wahhab MA. Antioxidant and antibacterial activity of silver nanoparticles biosynthesized using *Chenopodium murale* leaf extract. J Saudi Chem Soc. 2014;18:356-63. https://doi.org/10.1016/j. jscs.2013.09.011
- Harborne JB. Metode Fitokimia. 2nd ed. Indonesia: Penerbit ITB; 1987.
- 18. Kirom HS, Ramadhania ZM. Article Review: Biological Activity of *Acalypha indica* L. Farmaka. 2015;15(3):162-9.
- Komalamisra N, Trongtokit Y, Rongsriyam Y, Apiwathnasorn C. Screening for larvicidal activity in some Thai plants against four mosquito vector species. Southeast Asian J Trop Med Public Health. 2005;36(6):1412-2. https://doi.org/10.1002/ ptr.1637

PMid:16610643

 Kiran SR, Bhavani K, Devi PS, Rao BR, Reddy KJ. Composition and larvicidal activity of leaves and stem essential oils of Chloroxylon swietenia DC against Aedes aegypti and Anopheles stephensi. Bioresour Technol. 2006;97(18):2481-4. https://doi. org/10.1016/j.biortech.2005.10.003

PMid:16815011

- Isman MB. Botanical insecticides: For richer, for poorer. Pest Manag Sci. 2008;64(1):8-11. https://doi.org/10.1002/ps.1470 PMid:18022796
- Robinson T. Kandungan Organic Tumbuhan Tinggi. 6th ed. Indonesia: Penerbit ITB; 1995. p. 191-216.
- Wardani RS, Mifbakhuddin, Yokorinanti K. Effect of concentration of Tembelekan leaf extract (*Lantana camara*) on *Aedes aegypti larvae* death. J Kesehatan Masyarakat Indones. 2010;6(2):30-8. https://doi.org/10.33221/jikm.v9i01.820
- 24. Ahdiyah I, Purwani KI. The effect of the extract of the leaves of the Mangkokan (*Nothopanax scutellarium*) as the larvicide of *Culex* sp. Mosquito. 2015. [cited March 8, 2021]. Available online at http://webcache.googleusercontent.com/search?q=cache:XnJSEfBeYJ:ejurnal.its.ac.id/index.php/sains seni/article/view
- 25. Li J. Abrief Introduction to Citrus Limonoid. TAMU Colege.

- TAMUK Citrus Centre; 2021. Available from: http://www.ejaefche.wigo.es. [Last accessed on 2021 Mar 06].
- Diana L. Efektivitas Minyak Atsiri Kulit Jeruk Nipis (Citrus aurantifolia Swingle) Terhadap Mortalitas Larva Aedes aegypti L. Intisar III [Thesis]. Aceh: Universitas Syah Kuala; 2013. https://doi.org/10.31983/keslingmas.v37i2.3870
- 27. Pradani FY. Growth Index of Exposed Aedes aegypti L. Larvae in Water Extract of Jengkol (*Pithecellobium lobatum*) bark.
- Aspirator. 2009;1(2):81-5.
- 28. Gutierrez PM, Aubrey NA, Eugenio BA, Santos MF. Larvicidal activity of selected plant extracts against the dengue vector *Aedes aegypti* mosquito. Int Res J Biol Sci. 2014;3(4):23-32.
- Mardiana M, Supraptini S, Nunik SA. Datura metel Linnaeus as Botanical Insecticide and Larvicide and Raw Material for Traditional Medicine. Med Penelitian Pengembang Kesehatan. 2009;19(2):1-4.